

COMPARING Π_2^1 -PROBLEMS IN COMPUTABILITY THEORY AND REVERSE MATHEMATICS

DENIS R. HIRSCHFELDT

Reverse mathematics gives us a way to compare the relative strength of theorems by establishing implications and nonimplications over a weak subsystem of second-order arithmetic, typically RCA_0 , which corresponds roughly to computable mathematics. In many cases, nonimplications over RCA_0 are proved using ω -models, i.e., models of RCA_0 with standard first-order part. Implication over RCA_0 and over ω -models are not fine enough for some purposes, however, so other notions of computability-theoretic reduction between theorems have been extensively studied. These are particularly well-adapted to a class of theorems that includes a large proportion of those that have been studied in reverse mathematics: A Π_2^1 -*problem* is a sentence

$$\forall X [\Theta(X) \rightarrow \exists Y \Psi(X, Y)]$$

of second-order arithmetic such that Θ and Ψ are arithmetic. The term “problem” reflects a computability-theoretic view that sees such a sentence as a process of finding a suitable Y given an X satisfying certain conditions.

This talk will discuss some of these approaches to studying the relative strength of Π_1^2 -problems, focusing in particular on combinatorial examples, including work in [2, 1].

Acknowledgements. Hirschfeldt is partially supported by grants DMS-1600543 and DMS-1854279 from the National Science Foundation of the United States.

REFERENCES

- [1] D. D. Dzhabarov, D. R. Hirschfeldt, and S. C. Reitzes. Reduction games, provability, and compactness. In preparation.
- [2] D. R. Hirschfeldt and C. G. Jockusch, Jr. On notions of computability-theoretic reduction between Π_2^1 principles. *Journal of Mathematical Logic*, 16(1), 1650002, 59 pp., 2016.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, USA
E-mail address: `drh@math.uchicago.edu`